

Estimating Long-Term Effect of Siponimod on Disability Progression versus Virtual Placebo in SPMS Using RPSFT Model: EXPAND Data Up to 7 Years

Bruce A.C. Cree¹, Nicolas Rouyrre², Robert J. Fox³, Patrick Vermersch⁴, Gavin Giovannoni⁵, Amit Bar-Or⁶, Ralf Gold⁷, Jeff Maca⁸, Daniela Piani-Meier², Goeril Karlsson², Ludwig Kappos⁹

¹UCSF Weill Institute for Neurosciences, Department of Neurology, University of California San Francisco, San Francisco, CA, USA; ²Novartis Pharma AG, Basel, Switzerland; ³Mellen Center for Treatment and Research in Multiple Sclerosis, Neurological Institute, Cleveland, OH, USA; ⁴Univ. Lille, Inserm U1172 LiNCog, CHU Lille, FHU Precise, Lille, France; ⁵Blizard Institute, Barts and The London School of Medicine and Dentistry, Queen Mary University of London, London, United Kingdom; ⁶Center for Neuroinflammation and Experimental Therapeutics and Department of Neurology, Perelman School of Medicine, University of Pennsylvania, PA, USA; ⁷Department of Neurology, St Josef-Hospital/Ruhr-University Bochum, Bochum, Germany; ⁸Novartis Pharmaceuticals Corporation, East Hanover, NJ, USA; ⁹Research Center for Clinical Neuroimmunology and Neuroscience Basel (RC2NB), Departments of Medicine, Clinical Research, Biomedicine and Biomedical Engineering, University Hospital and University of Basel, Basel, Switzerland

Poster number: 004 (Neighborhood 4)

Session name: P6 (MS Clinical Trials and Therapeutics 2)

Session time: Sunday, April 3, 5:30 PM - 6:30 PM

Poster Presentation at the American Academy of Neurology (AAN) 2022, April 2-7, 2022



Scan to download a copy of this presentation

Author Name	Disclosures
Bruce A.C. Cree	Received personal compensation for consulting from Alexion, Atara, Autobahn, Avotres, Biogen, EMD Serono, Novartis, Sanofi, TG Therapeutics, and Therini. Received research support from Genentech.
Robert J. Fox	Received personal consulting fees from Actelion, Biogen, Celgene, EMD Serono, Genentech, Immunic, Novartis, and Teva. He has served on advisory committees for Actelion, Biogen, Immunic, and Novartis, and received clinical trial contract and research grant funding from Biogen and Novartis.
Patrick Vermersch	Received honoraria and consulting fees from Biogen Idec, Sanofi-Genzyme, Bayer, Novartis, Merck Serono, AB science, Imcysc, and Almirall, and research support from Biogen Idec, Sanofi-Genzyme, Bayer, and Merck Serono.
Gavin Giovannoni	Steering committee member on the daclizumab trials for AbbVie, the BG12 and daclizumab trials for Biogen, the fingolimod and siponimod trials for Novartis, the laquinimod trials for Teva and the ocrelizumab trials for Roche. He has also received consultancy fees for advisory board meetings for oral cladribine trials for Merck KGaA, Sanofi Genzyme, and in relation to DSMB activities for Synthon BV, as well as honoraria for speaking at the Physicians' summit and several medical education meetings. He is also the Co-Chief Editor of Multiple Sclerosis and Related Disorders (Elsevier).
Amit Bar-Or	Participated as a speaker in meetings sponsored by and received consulting fees and/or grant support from: Accure, Atara Biotherapeutics, Biogen, BMS/Celgene/Receptos, GlaxoSmithKline, Gossamer, Janssen/Actelion, Medimmune, Merck/EMD Serono, Novartis, Roche/Genentech, and Sanofi-Genzyme.
Ralf Gold	Received compensation for serving as a consultant or speaker from Bayer HealthCare, Biogen Idec, Merck Serono, Novartis, and Teva Neuroscience. He, or the institution he works for, has received research support from Bayer HealthCare, Biogen Idec, Merck Serono, Novartis, and Teva Neuroscience. He has also received honoraria as a Journal Editor from SAGE and Thieme Verlag.
Ludwig Kappos	Received the following exclusively for research support: Steering committee, advisory board, and consultancy fees from: Actelion, Bayer HealthCare, Biogen, BMS, Genzyme, Glaxo Smith Kline, Janssen, Japan Tobacco, Merck, Novartis, Roche, Sanofi, Santhera, Shionogi, TG Therapeutics; Speaker fees from: Bayer HealthCare, Biogen, Merck, Novartis, Roche, and Sanofi; Support of educational activities from: Allergan, Bayer HealthCare, Biogen, CSL Behring, Desitin, Genzyme, Merck, Novartis, Roche, Pfizer, Sanofi, Shire, and Teva; License fees for Neurostatus products and grants from: Bayer HealthCare, Biogen, European Union, InnoSwiss, Merck, Novartis, Roche, Swiss MS Society, and Swiss National Research Foundation.
Nicolas Rouyre	Employee of Novartis Gene Therapies, Inc., and owns Novartis stock or other equities.
Jeff Mecca	Employee of Novartis and may hold stocks of Novartis.
Daniela Piani-Meier, Goeril Karlsson	Employees of Novartis.

Funding source: This study is supported by Novartis Pharma AG, Basel, Switzerland.

Acknowledgments: Editorial support was provided by **Matthew Hartmann, PhD**, of Alphabet Health (New York, NY) and was funded by Novartis Pharmaceuticals Corporation. This poster was developed in accordance with Good Publication Practice (GPP3) guidelines. Authors had full control of the content and made the final decision on all aspects of this poster.

Introduction

- In the core part of the Phase 3 EXPAND study, siponimod significantly reduced the risk of 6-month confirmed disability progression (6mCDP) versus placebo by 26% in patients with secondary progressive multiple sclerosis (SPMS)¹
- In the long-term analysis (up to 5 years: data cut-off of April 2019) including data from the core and ongoing EXPAND-extension, the risk of 6mCDP was reduced by 22% and the time to 6mCDP was prolonged by 49% in the continuous siponimod group versus placebo-siponimod group²
- Long-term comparison of siponimod with placebo was not possible since placebo patients transitioned to open-label siponimod at the end of EXPAND core study (median duration of core part 21 months)
- Modelling of the long-term trajectory of patients initially randomized to placebo (virtual placebo) using Rank Preserving Structural Failure Time (RPSFT) was undertaken and then compared with long-term continuous siponimod group
 - In a previous analysis modelling a placebo treatment arm corrected for switch by RPSFT analysis in the overall SPMS population, the risk of 6mCDP in the continuous siponimod group was reduced by 31% (hazard ratio [HR]: 0.69, cut-off of April 2019) versus modelled placebo group and the time to 6mCDP was prolonged by ~50%–60%³
 - Different models were tested but only the RPSFT model produced predictions in a range consistent with the HR reported in the core part of the study. The accuracy and applicability of the RPSFT model to model a virtual placebo arm in the long term was further supported by simulations (for placebo patients after switching to siponimod) conducted under conditions similar to the EXPAND study, which included waning and increasing treatment effects³

Objective

- To present the long-term efficacy of continuous siponimod versus virtual placebo using the RPSFT model and uncorrected placebo-siponimod switch intent-to-treat (ITT) analysis in the overall SPMS population as well as in the active and non-active subgroups from the core and extension parts of the EXPAND study with up to 7 years of follow-up

1. Kappos L, et al. *Lancet* 2018;391:1263-1273. 2. Kappos L, et al. Oral presentation at AAN 2020. S40.003. 3. Cree BAC, et al. Poster Presented at the 8th Joint ACTRIMS-ECTRIMS Meeting, MSVirtual2020. P0013.

Introduction

Methods


Results

Conclusions



- The analysis included overall EXPAND SPMS population who received ≥ 1 dose of siponimod in the core study (ITT population)¹ and offered a switch to open-label siponimod in the ongoing extension (data cut-off: October 2020) for up to 7 years
- The long-term trajectory of placebo was estimated using the RPSFT model, a method that adjusts for treatment switch in trials with survival outcomes²⁻⁴
- In the present analysis, patients were grouped into the following:
 - Long-term continuous siponimod (LTCS) arm who received siponimod during the core and extension parts
 - Long-term placebo-siponimod arm (LTPS) who received placebo in the core and siponimod in the extension (ITT analysis, not modelled – ‘uncorrected’)
 - Long-term virtual placebo (LTVP) arm modelled (‘corrected’) by RPSFT method⁵

Study design



^aExtension data cut-off: Oct 2020 (Month 54 visit of extension); total study duration (core+extension): up to 7 years. ^bOpen-label starts when patient has an "event". EDSS, Expanded Disability Status Scale; EoCP, end of core part; ITT, intent-to-treat; LTCS, long-term continuous siponimod; LTPS, long-term placebo-siponimod arm; LTVP, long-term virtual placebo; MRI, magnetic resonance imaging; N, total number of patients (safety set); PYs, patient years; RPSFT, Rank Preserving Structural Failure Time; SPMS, secondary progressive multiple sclerosis; SDMT, Symbol Digit Modalities Test. 1. Kappos L, et al. *Lancet* 2018;391:1263-1273. 2. Latimer NR, et al. *Stat Methods Med Res.* 2017;26:724-751. 3. Latimer NR, et al. *Stat Methods Med Res.* 2018;27:765-784. 4. Korhonen P, et al. *J Biopharm Stat.* 2012;22:1258-1271. 5. Cree BAC, et al. Poster Presented at the 8th Joint ACTRIMS-ECTRIMS Meeting, MSVirtual2020. P0013.

- Data were analyzed in the overall population (siponimod [N=1105], placebo [N=546]) and in the subgroups of patients with active^a (siponimod [N=400], placebo-siponimod [N=182]) and non-active^b (siponimod [N=405], placebo-siponimod [N=207]) subgroups

Outcomes

- Time to 6mCDP based on the EDSS score in the continuous siponimod (LTCS) versus placebo-siponimod arm modelled (LTVP) and ITT analysis-not modelled (LTPS) arms


Statistical analysis

- For both ITT and RPSFT analysis, the time to 6mCDP was analyzed using a Cox proportional hazards model with treatment, country/region, baseline EDSS score and SPMS group (with-/without superimposed relapses, baseline definition) as covariates using combined EDSS data from the core and extension parts
- In the RPSFT analysis, patients randomized to placebo and crossing over to open-label siponimod were replaced by their counterfactual time to CDP without cross over based RPSFT model, e.g. modelling ('correcting') as if they had never switched to siponimod


^aDefined as the presence of at least one relapse in the 2 years before screening and/or ≥ 1 Gd+ T1 lesion at baseline. ^bDefined as no relapse in the 2 years prior to screening and no Gd+ T1 lesion at baseline. EDSS, Expanded Disability Status Scale; Gd+, gadolinium-enhancing; ITT, intent-to-treat; LTCS, long-term continuous siponimod; LTPS, long-term placebo-siponimod arm; LTVP, long-term virtual placebo; N, total number of patients (safety set); RPSFT, Rank Preserving Structural Failure Time; SPMS, secondary progressive multiple sclerosis.

- **Overall population:** The risk of 6mCDP was reduced by 33% ($p < 0.0001$) and the median time to 6mCDP was prolonged by 62%
- **Active SPMS population:** The risk of 6mCDP was reduced by 42% ($p < 0.0001$) and the median time to 6mCDP was prolonged by 79%
- **Non-active SPMS population:** The risk of 6mCDP was reduced by 20% ($p = 0.0534$) and the median time to 6mCDP was prolonged by 44%


Overall SPMS population



Active SPMS subgroup



Non-active SPMS subgroup



*Indicates statistical significance (2-sided) at the 0.05 level

6mCDP, 6-month confirmed disability progression; CI, confidence interval; HR, hazard ratio; ITT, intent-to-treat; LTCS, long-term continuous siponimod; LTVP, long-term placebo-siponimod (ITT analysis-not modelled); LTVP, long-term modelled virtual placebo; RPSFT, Rank Preserving Structural Failure Time; SPMS, secondary progressive multiple sclerosis

Introduction

Methods

Results

Conclusions



- This analysis confirmed the sustained efficacy of continuous siponimod up to 7 years in significantly reducing the risk of progression and prolonging the time to 6mCDP versus both modelled virtual placebo and placebo-siponimod switch ITT analysis (not modelled placebo-siponimod arm) in the overall SPMS patients
- In the active SPMS subgroup, the effect of siponimod on disability progression was more pronounced and sustained for up to 7 years
- In the non-active SPMS subgroup, a sustained strong trend favoring continuous siponimod was observed; longer time to 6mCDP in the placebo arm compared with the placebo arm in the active SPMS subgroup suggests longer observation periods are required to uncover the full impact of treatment in non-active SPMS subgroup
- RPSFT virtual placebo arm can be used to estimate the long-term treatment benefits of siponimod in reducing disability progression
- One limitation of this analysis is that it assumes an attrition (before observing the 6mCDP) which is balanced between treatment arms. Given that all patients are currently receiving open-label siponimod, this assumption is reasonable

6mCDP, 6-month confirmed disability progression; RPSFT, Rank Preserving Structural Failure Time; SPMS, secondary progressive multiple sclerosis

Introduction

Methods

Results

Conclusions

